arXiv:1006.0074v1 [math.CA] 1 Jun 2010

SECOND–ORDER LINEAR CONSTANT COEFFICIENT DYNAMIC EQUATIONS WITH POLYNOMIAL FORCING ON TIME SCALES

DOUGLAS R. ANDERSON

Abstract. A general solution for a second-order linear constant coefficient dynamic equation with polynomial forcing on time scales is given.

1. POLYNOMIAL FORCING OF SECOND-ORDER ORDINARY DYNAMIC EQUATIONS WITH CONSTANT COEFFICIENTS

Recall that polynomials on time scales [1] are defined via $h_0(t,a) \equiv 1$ and recursively for $k \in \mathbb{N}$ by

$$h_k(t,a) = \int_a^t h_{k-1}(\tau,a)\Delta\tau.$$

We have the following result.

Theorem 1.1 (Constant Coefficients with Polynomial Forcing). *Let $a \in \mathbb{T}$ and $\alpha, \beta \in \mathbb{R}$. A general solution of the second-order linear constant coefficient equation with polynomial forcing*

$$y^{\Delta\Delta} + 2\alpha y^{\Delta} + \beta y = \sum_{i=0}^{k} \gamma_i h_i(\cdot,a), \quad \gamma_i \in \mathbb{R}, \tag{1.1}$$

with $\beta \neq 0$, $\beta \neq \alpha^2$, and $1 - 2\alpha\mu + \beta\mu^2 \neq 0$ is given by

$$y(t) = c_1 e_{\lambda_1}(t,a) + c_2 e_{\lambda_2}(t,a) + \sum_{i=0}^{k} \xi_i h_i(t,a), \tag{1.2}$$

where $\lambda_1 = -\alpha - \sqrt{\alpha^2 - \beta}$, $\lambda_2 = -\alpha + \sqrt{\alpha^2 - \beta}$, $\xi_{k-1} = \frac{1}{\beta}\left(\gamma_{k-1} - \frac{2\alpha}{\beta}\gamma_k\right)$, $\xi_k = \frac{1}{\beta}\gamma_k$,

$$\xi_i = \omega_1 \lambda_1^i + \omega_2 \lambda_2^i + \sum_{\tau=0}^{i-1} \sum_{s=0}^{\tau-1} \gamma_s \lambda_1^{i+s-2\tau} \beta^{\tau-1-s} \quad \textit{for} \quad i \in \{0, \cdots, k-2\},$$

2000 *Mathematics Subject Classification.* 34N05, 26E70, 39A10.

Key words and phrases. Ordinary difference equations; ordinary dynamic equations; inhomogeneous equations; time scales; reduction of order.

and

$$\begin{bmatrix}\omega_1\\ \omega_2\end{bmatrix} = \frac{1}{2\beta^{k-1}\sqrt{\alpha^2-\beta}}\begin{bmatrix}\lambda_2^k & -\lambda_2^{k-1}\\ -\lambda_1^k & \lambda_1^{k-1}\end{bmatrix}\begin{bmatrix}\frac{1}{\beta}\left(\gamma_{k-1}-\frac{2\alpha}{\beta}\gamma_k\right)-\sum_{\tau=0}^{k-2}\sum_{s=0}^{\tau-1}\gamma_s\lambda_1^{k-1+s-2\tau}\beta^{\tau-1-s}\\ \frac{1}{\beta}\gamma_k-\sum_{\tau=0}^{k-1}\sum_{s=0}^{\tau-1}\gamma_s\lambda_1^{k+s-2\tau}\beta^{\tau-1-s}\end{bmatrix}.$$

Proof. By the theory of linear dynamic equations [3, Theorem 3.16], a general solution of (1.1) has the form

$$y(t) = c_1e_{\lambda_1}(t,a) + c_2e_{\lambda_2}(t,a) + y_d(t) \tag{1.3}$$

for

$$\lambda_1 = -\alpha - \sqrt{\alpha^2-\beta} \quad \text{and} \quad \lambda_2 = -\alpha + \sqrt{\alpha^2-\beta}, \tag{1.4}$$

where the time scale exponential functions $e_{\lambda_1}(\cdot,a)$ and $e_{\lambda_2}(\cdot,a)$ are solutions of the corresponding homogeneous equation

$$y^{\Delta\Delta} + 2\alpha y^{\Delta} + \beta y = 0,$$

and y_d is a particular solution of the inhomogeneous equation (1.1). Here $y_1 = e_{\lambda_1}(\cdot,a)$ and $y_2 = e_{\lambda_2}(\cdot,a)$ are linearly independent since their Wronskian

$$W(y_1,y_2) = y_1y_2^{\Delta} - y_1^{\Delta}y_2 = (\lambda_2-\lambda_1)y_1y_2 = 2\sqrt{\alpha^2-\beta}\, e_{\lambda_1\oplus\lambda_2}(\cdot,a) = 2\sqrt{\alpha^2-\beta}\, e_{(-2\alpha+\mu\beta)}(\cdot,a)$$

is well defined due to the regressivity assumption $1-2\alpha\mu+\beta\mu^2 \neq 0$ and is never zero on $\mathbb{T}$. We guess that a particular solution y_d has the form

$$y_d = \sum_{i=0}^{k}\xi_ih_i(\cdot,a), \qquad \xi_i \in \mathbb{R}.$$

Then

$$y_d^{\Delta} = \sum_{i=0}^{k-1}\xi_{i+1}h_i(\cdot,a) \quad \text{and} \quad y_d^{\Delta\Delta} = \sum_{i=0}^{k-2}\xi_{i+2}h_i(\cdot,a).$$

Plugging into (1.1) and solving for the coefficients we have

$$\xi_{k-1} = \frac{1}{\beta}\left(\gamma_{k-1}-\frac{2\alpha}{\beta}\gamma_k\right) \quad \text{and} \quad \xi_k = \frac{1}{\beta}\gamma_k, \tag{1.5}$$

while the rest of the coefficients ξ_i satisfy the second-order linear constant coefficient difference equation with forcing given by

$$\xi_{i+2} + 2\alpha\xi_{i+1} + \beta\xi_i = \gamma_i, \quad i = 0,\dots,k-2. \tag{1.6}$$

The preceding difference equation (1.6) has the solution [2, Section 4]

$$\xi_i = \omega_1\lambda_1^i + \omega_2\lambda_2^i + \sum_{\tau=0}^{i-1}\sum_{s=0}^{\tau-1}\gamma_s\lambda_1^{i+s-2\tau}\beta^{\tau-1-s}$$

where λ_1 and λ_2 are given in (1.4) above. Using the terminal conditions in (1.5) and simple linear algebra we have

$$\begin{bmatrix}\omega_1\\ \omega_2\end{bmatrix} = \frac{1}{2\beta^{k-1}\sqrt{\alpha^2-\beta}}\begin{bmatrix}\lambda_2^k & -\lambda_2^{k-1}\\ -\lambda_1^k & \lambda_1^{k-1}\end{bmatrix}\begin{bmatrix}\frac{1}{\beta}\left(\gamma_{k-1}-\frac{2\alpha}{\beta}\gamma_k\right)-\sum_{\tau=0}^{k-2}\sum_{s=0}^{\tau-1}\gamma_s\lambda_1^{k-1+s-2\tau}\beta^{\tau-1-s}\\ \frac{1}{\beta}\gamma_k-\sum_{\tau=0}^{k-1}\sum_{s=0}^{\tau-1}\gamma_s\lambda_1^{k+s-2\tau}\beta^{\tau-1-s}\end{bmatrix}.$$

□

References

[1] R. P. Agarwal and M. Bohner, Basic calculus on time scales and some of its applications, *Resultate der Mathematik*, 35 (1-2): 3–22 (1999).

[2] D. R. Anderson and C. C. Tisdell, Alternative solutions of inhomogeneous second-order linear dynamic equations on time scales, *Journal of Difference Equations and Applications*, to appear.

[3] M. Bohner and A. Peterson, *Dynamic Equations on Time Scales, An Introduction with Applications*, Birkhäuser, Boston, 2001.

[4] S. Hilger, Analysis on measure chains – a unified approach to continuous and discrete calculus, *Results Math.*, 18 (1990) 18–56.

Department of Mathematics and Computer Science, Concordia College, Moorhead, MN 56562 USA, visiting United International College, Zhuhai, Guangdong, China

E-mail address: `andersod@cord.edu`

URL: `http://www.cord.edu/faculty/andersod/bib.html`